


# Glycoprotein 36 Antibody / Gp36 / Podoplanin [clone PDPN/8875R] (V4701)

Catalog No.	Formulation	Size
V4701-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4701-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4701SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


## Recombinant RABBIT MONOCLONAL

#### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	PDPN/8875R
Purity	Protein A/G affinity
UniProt	Q86YL7
Localization	Cell surface
Applications	ELISA (Order BSA-free Format For Coating) : Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Glycoprotein 36 antibody is available for research use only.



IHC staining of FFPE human prostate tissue with Glycoprotein 36 antibody (clone PDPN/8875R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



## **Description**

It recognizes a muco-protein of 38-43kDa, which is identified as Podoplanin (PDPN), also called Glycoprotein 36 (Gp36). It localizes in stromal cells of peripheral lymphoid tissue and thymic epithelial cells. As a regulator of the lymphatic endothelium, podoplanin probably plays a role in maintaining the unique shape of podocytes. It is selectively expressed in lymphatic endothelium as well as lymphoangiomas, Kaposi sarcomas, and in a subset of angiosarcomas with probable lymphatic differentiation. Recent studies have also shown podoplanin to be a highly sensitive and relatively specific marker for epithelioid mesothelioma. Therefore, it can be used in a panel to distinguish mesotheliomas or mesothelial cells from pulmonary carcinomas.

## **Application Notes**

Optimal dilution of the Glycoprotein 36 antibody should be determined by the researcher.

#### **Immunogen**

A recombinant partial protein sequence (within amino acids 1-150) from the human protein was used as the immunogen for the Glycoprotein 36 antibody.

## **Storage**

Aliquot the Glycoprotein 36 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.