
Glypican 3 Antibody GPC3/7107 / Mouse Monoclonal [clone GPC3/7107] (V4020)


Catalog No.	Formulation	Size
V4020-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4020-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4020SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	GPC3/7107
Purity	Protein A affinity
UniProt	P51654
Applications	Immunohistochemistry (FFPE): 1-2ug/ml
Limitations	This Glypican 3 Antibody GPC3/7107 is available for research use only.



IHC staining of FFPE human prostate tissue with Glypican 3 Antibody GPC3/7107. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Glypican 3 antibody (clone GPC3/7107) as confirmation of integrity and purity.

Description

Glypican-3 (GPC3) is a glycosylphospatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilm s tumor. In patients with HCC, GPC3 is overexpressed in neoplastic liver tissue and elevated in serum, but is undetectable in normal liver, benign liver, and the serum of healthy donors. GPC3 expression is also found to be higher in HCC liver tissue than in cirrhotic liver or liver with focal lesions such as dysplastic nodules and areas of hepatic adenoma (HA) with malignant transformation. In the context of testicular germ cell tumors, GPC3 expression is up regulated in certain histologic subtypes, specifically yolk sac tumors and choriocarcinoma. A high level of GPC3 expression is also found in some types of embryonal tumors, such as Wilm s tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.

For an alternative monoclonal antibody targeting this cell-surface proteoglycan, see our GPC3 Antibody rGPC3/863, validated for immunohistochemistry, immunofluorescence, and flow cytometry.

Application Notes

Optimal dilution of the Glypican 3 Antibody GPC3/7107 should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 from the human GPC3 protein was used as the immunogen for the Glypican 3 antibody.

Storage

Aliquot the Glypican 3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.