


Granulocyte Marker Antibody [clone BM-2] (V2339)

Catalog No.	Formulation	Size
V2339-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2339-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2339SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2339IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


 Citations (1)

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	BM-2
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
Gene ID	Unknown
Localization	Cytoplasmic
Applications	Flow Cytometry : 0.5-1ug/10 ⁶ cells Immunofluorescence : 1-2ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This Granulocyte Marker antibody is available for research use only.



IHC staining of tonsil tissue with Granulocyte Marker antibody (clone BM-2). Note specific cytoplasmic staining.



SDS-PAGE Analysis of Purified, BSA-Free Granulocyte Marker Antibody (clone BM-2). Confirmation of Integrity and Purity of the Antibody.

Description

This antibody recognizes an unidentified antigen in the cytoplasm of mature granulocytes and can be used as a granulocyte marker in normal tissues or inflammatory processes. The granulocyte marker antibody shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. Clone BM-2 mAb reacts with early precursor and mature forms of human myeloid cells. It is useful for the detection of myeloid leukemias and granulocytic sarcomas.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the Granulocyte Marker antibody to be titered up or down for optimal performance.

1. No special pretreatment is required for staining of formalin/paraffin tissues.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Nuclei from pokeweed mitogen stimulated human peripheral blood lymphocytes were used as the immunogen for this Granulocyte Marker antibody.

Storage

Store the Granulocyte Marker antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

References (1)