


H-FABP Antibody / Heart type fatty acid binding protein / FABP3 [clone FABP3/3770] (V5694)


Catalog No.	Formulation	Size
V5694-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5694-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5694SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	FABP3/3770
Purity	Protein G affinity
UniProt	P05413
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 2-4ug/ml
Limitations	This H-FABP antibody is available for research use only.



H-FABP Antibody Human Cardiac Muscle Immunohistochemistry. IHC staining of FFPE human heart tissue with PSAP antibody (clone FABP3/3770). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



H-FABP Antibody Human Cardiac Muscle WB. Western blot testing of human heart tissue lysate with H-FABP antibody (clone FABP3/3770). Predicted molecular weight ~14 kDa.

Description

The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is a candidate tumor suppressor gene for human breast cancer. Alternative splicing results in multiple transcript variants.

Explore our [FABP3 Antibody page](#) for additional western blot, immunohistochemistry, and microarray specificity validation data supporting studies of myocardial lipid transport and oxidative metabolism.

Application Notes

Optimal dilution of the H-FABP antibody should be determined by the researcher.

Immunogen

A human partial recombinant protein (from within amino acids 1-127) was used as the immunogen for the H-FABP antibody.

Storage

Aliquot the H-FABP antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.