


HA Antibody (F52062)


Catalog No.	Formulation	Size
F52062-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F52062-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	HA tag
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
Applications	Western Blot : 1:1000 Immunofluorescence : 1:10-1:50 ChIP : 1:100 Immunoprecipitation : suitable
Limitations	This HA antibody is available for research use only.



Western blot analysis of lysate from 12 tagged recombinant protein cell lines using HA antibody diluted at 1:1000.



Western blot analysis of HA antibody and tagged recombinant protein bacterial lysate



GTPase activity of Dynamin-2 is required for endocytosis of cell-surface tTG. MRC-5 fibroblasts were transiently transfected with either wild-type (wt) or a GTPase-deficient Dynamin-2 mutant with a hemagglutinin (HA) tag.(Provided by Evgeny A. Zemskov, Irina Mikhailenko)

Description

Epitope tags consisting of short sequences recognized by well-characterized antibodies have been widely used in the study of protein expression in various systems. The HA tag (YPYDVPDYAYPYDVPDYA) is an established example. NSJBio's anti-HA polyclonal antibody provides a simple solution to detect the expression of an HA-tagged protein in cells.

Application Notes

Titration of the HA antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

The sequence CYPYDVPDYAYPYDVPDYA was used as the immunogen for this HA antibody.

Storage

Aliquot the HA antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.