



# HA Antibody (F52062)


Catalog No.	Formulation	Size
F52062-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F52062-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	HA tag
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
Applications	Western Blot : 1:1000 Immunofluorescence : 1:10-1:50 ChIP : 1:100 Immunoprecipitation : suitable
Limitations	This HA antibody is available for research use only.







GTPase activity of Dynamin-2 is required for endocytosis of cell-surface tTG. MRC-5 fibroblasts were transiently transfected with either wild-type (wt) or a GTPase-deficient Dynamin-2 mutant with a hemagglutinin (HA) tag.(Provided by Evgeny A. Zemskov, Irina Mikhailenko)

### **Description**

Epitope tags consisting of short sequences recognized by well-characterizated antibodies have been widely used in the study of protein expression in various systems. The HA tag (YPYDVPDYAYPYDVPDYA) is an established example. NSJBio's anti-HA polyclonal antibody provides a simple solution to detect the expression of an HA-tagged protein in cells.

### **Application Notes**

Titration of the HA antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

The sequence CYPYDVPDYAYPYDVPDYA was used as the immunogen for this HA antibody.

#### **Storage**

Aliquot the HA antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.