
HCG-beta Antibody / HCGb [clone HCGb/7201R] (V4001)

Catalog No.	Formulation	Size
V4001-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4001-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4001SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	HCGb/7201R
Purity	Protein A/G affinity
UniProt	P01233
Localization	Cytoplasm, secreted
Applications	Immunohistochemistry (FFPE): 1-2ug/ml
Limitations	This HCG-beta antibody is available for research use only.



HCG-beta Antibody Human Pituitary Gland Tissue IHC. Immunohistochemistry staining of FFPE human pituitary gland tissue with HCG-beta antibody (clone HCGb/7201R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



HCG-beta Antibody Human Placenta IHC. Immunohistochemistry staining of FFPE human placental tissue with HCG-beta antibody (clone HCGb/7201R). Negative control inset: PBS used instead of primary antibody to control for secondary Ab binding.

Description

Human chorionic gonadotropin antibody (hCG) is a glycoprotein hormone synthesized in syncytiotrophoblastic cells of placenta and in certain trophoblastic tumors. The hormone-specific alpha chains have molecular weights of 13 kDa. HCG is found in moles and choriocarcinoma, chorionic components of germ cell tumors, and syncytiotrophoblast like cells in seminoma/dysgerminoma and embryonal carcinoma. In diagnostic pathology, hCG is a useful marker for classification of germ cell tumors, identification of extragonadal germ cell tumors.

This antibody is part of a broader antibody panel offered by NSJ Bioreagents.

Application Notes

Optimal dilution of the HCG-beta antibody should be determined by the researcher.

Immunogen

Recombinant full-length human protein was used as the immunogen for the HCG-beta antibody.

Storage

Aliquot the HCG-beta antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.