


HCG-beta Antibody [clone HCGb/211] (V2374)

Catalog No.	Formulation	Size
V2374-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2374-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2374SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	HCGb/211
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
Gene ID	1082
Localization	Cytoplasmic, Secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This HCG-beta antibody is available for research use only.



SDS-PAGE Analysis of Purified, BSA-Free HCG-beta Antibody (clone HCGb/211). Confirmation of Integrity and Purity of the Antibody.

Description

This antibody reacts with a protein of 22kDa, identified as beta subunit of HCG. It does not cross react with the alpha subunit. HCG is a glycoprotein which is secreted in large quantities by normal trophoblasts. It is present only in trace

amounts in non-pregnant urine and sera but rises sharply during pregnancy. HCG is composed of two non-identical, non-covalently linked polypeptide chains designated as the alpha and beta subunits. The alpha subunit is identical to that of thyroid stimulating hormone (TSH), follicle stimulating hormone (FSH), and luteinizing hormone (LH). HCG-beta antibody detects cells and tumors of trophoblastic origin such as choriocarcinoma. Large cell carcinoma and adenocarcinoma of the lung demonstrate antibody positivity in 90% and 60% of cases respectively; 20% of lung squamous cell carcinomas are positive. HCG expression by non-trophoblastic tumors may indicate aggressive behavior.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the HCG-beta antibody to be titrated up or down for optimal performance.

1. Clone HCGb/211 (recommended capture) will pair with clones [HCGb/459](#) and [HCGb/54](#) (recommended detect) by ELISA.

Immunogen

Purified HCG-beta protein was used as the immunogen for this antibody.

Storage

Store the HCG-beta antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Alternate Names

CG-beta; CGB3; CGB5; CGB7; CGB8; Choriogonadotropin Subunit beta; hCGB, hCG-beta antibody

References (1)