


## HDAC6 Antibody [clone PCR-P-HDAC6-1A4] (V9524)


Catalog No.	Formulation	Size
V9524-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9524-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9524SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a
<b>Clone Name</b>	PCR-P-HDAC6-1A4
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q9UBN7
<b>Localization</b>	Nuclear
<b>Applications</b>	Flow Cytometry : 1-2ug/million cells Immunofluorescence : 1-2ug/ml
<b>Limitations</b>	This HDAC6 antibody is available for research use only.



Immunofluorescent staining of PFA-fixed human HeLa cells using HDAC6 antibody (green, clone PCR-P-HDAC6-1A4) and phalloidin (red).




FACS staining of PFA-fixed human HeLa cells with HDAC6 antibody (blue, clone PCRP-HDAC6-1A4) and unstained cells (red).



SDS-PAGE analysis of purified, BSA-free HDAC6 antibody (clone PCRP-HDAC6-1A4) as confirmation of integrity and purity.

#### Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using HDAC6 antibody (clone PCRP-HDAC6-1A4). These results demonstrate the foremost specificity of the PCRP-HDAC6-1A4 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

## Description

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, PCAF (p300/CBP associated factor), p300/CBP, HAT1, and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated RPD3) and HDAC3-6, have been identified as histone deacetylases. This enzyme deacetylates lysine residues in histones H2A, H2B, H3 and H4.

## Application Notes

Optimal dilution of the HDAC6 antibody should be determined by the researcher.

## Immunogen

Recombinant full-length human HDAC6 protein was used as the immunogen for the HDAC6 antibody.

## Storage

Aliquot the HDAC6 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

