


## Heart Fatty Acid Binding Protein Antibody / FABP3 [clone FABP3/8343R] (V4876)

Catalog No.	Formulation	Size
V4876-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4876-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4876SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG, kappa
<b>Clone Name</b>	FABP3/8343R
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P05413
<b>Localization</b>	Cytoplasm
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This Heart Fatty Acid Binding Protein antibody is available for research use only.



Heart Fatty Acid Binding Protein Antibody Heart Tissue Immunohistochemistry. IHC staining of FFPE human heart tissue with Heart Fatty Acid Binding Protein antibody (clone FABP3/8343R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

### Description

The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three

distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is a candidate tumor suppressor gene for human breast cancer. Alternative splicing results in multiple transcript variants.

Explore our [FABP3 Antibody page](#) for additional western blot, immunohistochemistry, and microarray specificity validation data supporting studies of myocardial lipid transport and oxidative metabolism.

## Application Notes

Optimal dilution of the Heart Fatty Acid Binding Protein antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 1-127) from the human protein was used as the immunogen for the Heart Fatty Acid Binding Protein antibody.

## Storage

Aliquot the Heart Fatty Acid Binding Protein antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.