
# Helicobacter pylori Antibody (Catalase) [clone HPYL/7226] (V9762)


| Catalog No. | Formulation | Size |
|---|---|---|
| V9762-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V9762-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V9762SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Helicobacter pylori |
| **Format** | Purified |
| **Host** | Mouse |
| **Clonality** | Monoclonal (mouse origin) |
| **Isotype** | Mouse IgG2b, kappa |
| **Clone Name** | HPYL/7226 |
| **Purity** | Protein A/G affinity |
| **UniProt** | Not Applicable |
| **Localization** | Cell surface, Cytoplasm |
| **Applications** | Immunohistochemistry (FFPE) : 1-2ug/ml |
| **Limitations** | This Helicobacter pylori antibody is available for research use only. |



IHC staining of H. pylori-infected FFPE human stomach tissue with Helicobacter pylori antibody (clone HPYL/7226). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE H. pylori-infected human stomach tissue with Helicobacter pylori antibody (clone HPYL/7226). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

The spiral shaped bacterium Helicobacter pylori is strongly associated with inflammation of the stomach and is also implicated in the development of gastric malignancy. H. pylori is known to cause peptic ulcers and chronic gastritis in human. It is associated with duodenal ulcers and may be involved in development of adenocarcinoma and low-grade lymphoma of mucosa associated lymphoid tissue in the stomach. This antibody stains the individual H. pylori bacterium when it presents on the surface of the epithelium or in the cytoplasm of the epithelial cells in biopsy tissue sections from the antrum and body of the stomach.

## Application Notes

Optimal dilution of the Helicobacter pylori antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 323-445 from the Helicobacter pylori Catalase protein was used as the immunogen for the Helicobacter pylori antibody.

## Storage

Aliquot the Helicobacter pylori antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.


Copyright © NSJ Bioreagents. All rights reserved.