


HER-4 Antibody / ERBB4 [clone HFR-1] (V7741)


Catalog No.	Formulation	Size
V7741-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7741-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7741SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	HFR-1
Purity	Protein G affinity chromatography
UniProt	Q15303
Localization	Cytoplasmic, plasma membrane, nuclear
Applications	Flow Cytometry : 1-2ug/10^6 cells in 0.1ml Immunofluorescence : 1-2ug/ml
Limitations	This HER-4 antibody is available for research use only.



Immunofluorescent staining of permeabilized human MCF7 cells with HER-4 antibody (clone HFR-1, green) and Reddot nuclear stain (red).



Flow cytometry testing of PFA-fixed human MCF7 cells with HER-4 antibody (clone HFR-1); Red=isotype control, Blue= HER-4 antibody.



SDS-PAGE analysis of purified, BSA-free HER-4 antibody (clone HFR-1) as confirmation of integrity and purity.

Description

The EGF receptor family comprises several related receptor tyrosine kinases that are frequently overexpressed in a variety of carcinomas. Members of this receptor family include EGFR (HER1), Neu (ErbB-2, HER2), ErbB-3 (HER3) and ErbB-4 (HER4), which form either homodimers or heterodimers upon ligand binding. The gene encoding ErbB-4 is expressed as a full-length protein, which produces a short membrane-anchored cytoplasmic domain fragment and a long ectodomain fragment. The short fragment is heavily tyrosine phosphorylated and possesses tyrosine kinase catalytic activity toward an exogenous substrate. Proteolytic cleavage of ErbB-4 is promoted by the binding of heregulin. ErbB-4 is involved in cell proliferation and differentiation and its expression is highest in breast carcinoma cell lines, normal skeletal muscle, heart, pituitary, brain and cerebellum. Its expression in breast cancer, pediatric brain cancer and other types of carcinomas has been reported in studies which suggest ErbB4 expression is involved in both normal tissue development and carcinogenesis.

Application Notes

Optimal dilution of the HER-4 antibody should be determined by the researcher.

Immunogen

A recombinant human partial protein (amino acids 1116-1269) was used as the immunogen for the HER-4 antibody.

Storage

Store the HER-4 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).