


# Hexokinase 1 Antibody / HK1 (RQ5501)


Catalog No.	Formulation	Size
RQ5501	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P19367
Applications	Western Blot: 0.25-0.5ug/ml Immunohistochemistry (FFPE): 1-2ug/ml Immunofluorescence: 2-4ug/ml Direct ELISA: 0.1-0.5ug/ml
Limitations	This Hexokinase 1 antibody is available for research use only.




IHC staining of FFPE human rectal cancer with Hexokinase 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer with Hexokinase 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse brain with Hexokinase 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat brain with Hexokinase 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human U-2 OS cells with Hexokinase 1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human 1) placenta, 2) U-2 OS, 3) HEK293, 4) A549 and rat 5) brain, 6) heart, 7) lung and mouse 8) brain, 9) heart and 10) lung lysate with Hexokinase 1 antibody. Predicted molecular weight ~102 kDa.

## **Description**

Hexokinase-1 (HK1) is an enzyme that in humans is encoded by the HK1 gene on chromosome 10. It is mapped to 10q22.1. Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in several transcript variants which encode different isoforms, some of which are tissue-specific.

# **Application Notes**

Optimal dilution of the Hexokinase 1 antibody should be determined by the researcher.

### **Immunogen**

A human recombinant protein (amino acids D17-R323) was used as the immunogen for the Hexokinase 1 antibody.

#### **Storage**

After reconstitution, the Hexokinase 1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store

