


# Hexokinase II Antibody / HK2 (R32605)


Catalog No.	Formulation	Size
R32605	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

### **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P52789
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This Hexokinase II antibody is available for research use only.




Western blot testing of human 1) U-87 MG, 2) HEL, 3) HepG2 and 4) RT4 cell lysate with Hexokinase II antibody at 0.5ug/ml. Predicted molecular weight ~102 kDa.




Western blot testing of 1) human HeLa, 2) human MCF7, 3) human HepG2, 4) rat skeletal muscle, 5) rat heart, 6) mouse skeletal muscle and 7) mouse heart tissue lysate with Hexokinase II antibody at 0.5ug/ml. Predicted molecular weight ~102 kDa.




IHC staining of FFPE human breast cancer tissue with Hexokinase II antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE mouse skeletal muscle tissue with Hexokinase II antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat skeletal muscle tissue with Hexokinase II antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Flow cytometry testing of fixed and permeabilized human HepG2 cells with Hexokinase II antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Hexokinase II antibody.

## **Description**

Hexokinase II, also known as HK2, is an enzyme which in humans is encoded by the HK2 gene on chromosome 2. Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes hexokinase ii/2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate of glycolysis seen in rapidly growing cancer cells.

### **Application Notes**

Optimal dilution of the Hexokinase II antibody should be determined by the researcher.

#### **Immunogen**

Amino acids 460-497 (AYRLADQHRARQKTLEHLQLSHDQLLEVKRRMKVEMER-human) were used as the immunogen for the Hexokinase II antibody.

#### **Storage**

After reconstitution, the Hexokinase II antibody can be stored for up to one month at 4oC. For long-term, aliquot and store

