

## HIF1 alpha Antibody (F48865)


Catalog No.	Formulation	Size
F48865-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F48865-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Hamster
<b>Predicted Reactivity</b>	Bovine
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Purified
<b>UniProt</b>	Q16665
<b>Localization</b>	Nuclear, possible cytoplasmic
<b>Applications</b>	Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50 Immunofluorescence : 1:10-1:50
<b>Limitations</b>	This HIF1 alpha antibody is available for research use only.

250  
130  
95  
72  
55

Western blot analysis of HIF1 alpha antibody and CHO lysate. Routinely observed molecular weight: 100~120 kDa.



HIF1 alpha antibody IHC analysis in formalin fixed and paraffin embedded lung carcinoma.



Fluorescent confocal image of HeLa cell stained with HIF1 alpha antibody at 1:25. HIF1A immunoreactivity is localized to the cytoplasm and nucleus.

## Description

Hypoxia-inducible factor-1 (HIF1) is a transcription factor found in mammalian cells cultured under reduced oxygen tension that plays an essential role in cellular and systemic homeostatic responses to hypoxia. HIF1 is a heterodimer composed of an alpha subunit and a beta subunit. The beta subunit has been identified as the aryl hydrocarbon receptor nuclear translocator (ARNT). This protein encodes the alpha subunit of HIF-1. Overexpression of a natural antisense transcript (aHIF) of this gene has been shown to be associated with nonpapillary renal carcinomas.

## Application Notes

Titration of the HIF-1 alpha antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 1-30 from the human protein was used as the immunogen for this HIF1 alpha antibody.

## Storage

Aliquot the HIF1 alpha antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.