


# **Histone H2AX Antibody (RQ5283)**


Catalog No.	Formulation	Size
RQ5283	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P16104
Applications	Western Blot: 0.5-1ug/ml Immunohistochemistry (FFPE): 2-5ug/ml Flow Cytometry: 1-3ug/million cells Immunoprecipitation: 2ug/500ug of lysate Direct ELISA: 0.1-0.5ug/ml
Limitations	This Histone H2AX antibody is available for research use only.




IHC staining of FFPE human breast cancer tissue with Histone H2AX antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse liver tissue with Histone H2AX antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human HEL, 2) human PC-3, 3) human Caco-2, 4) rat PC-12, 5) rat C6 and 6) mouse NIH 3T3 cell lysate with Histone H2AX antibody. Predicted molecular weight ~15 kDa.



Flow cytometry testing of fixed and permeabilized human PC-3 cells with Histone H2AX antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Histone H2AX antibody.



Immunoprecipitation of Histone H2AX protein from 500ug of human 293T whole cell lysate with 2ug of Histone H2AX antibody.

## **Description**

H2A histone family member X (usually abbreviated as H2AX) is a type of histone protein from theH2A family encoded by the H2AFX gene. Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

## **Application Notes**

Optimal dilution of the Histone H2AX antibody should be determined by the researcher.

#### **Immunogen**

Amino acids S2-T121 from the human protein were used as the immunogen for the Histone H2AX antibody.

## **Storage**

Store the Histone H2AX antibody at -20oC.