

## HLA-DQ Antibody (MHC II) [clone SPM422] (V2582)

Catalog No.	Formulation	Size
V2582-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2582-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2582SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

### Bulk quote request

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a, kappa
<b>Clone Name</b>	SPM422
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P01908, P01909, P01920
<b>Localization</b>	Cell surface
<b>Applications</b>	ELISA (order BSA/sodium Azide-free Format For Coating) : Flow Cytometry : 0.5-1ug/10 <sup>6</sup> cells Immunofluorescence : 0.5-1ug/ml
<b>Limitations</b>	This HLA-DQ antibody is available for research use only.



## Description

Recognizes a DQ antigen, which is a dimer of 60kDa. The class II molecule is a heterodimer consisting of an alpha (DQA) and a beta chain (DQB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B Lymphocytes, dendritic cells, macrophages). The alpha chain is approximately 33-35kDa. It is encoded by 5 exons; exon 1 encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. Within the DQ molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to four different molecules. Typing for these polymorphisms is routinely done for bone marrow transplantation.

## Application Notes

Optimal dilution of the HLA-DQ antibody should be determined by the researcher.

1. No special pretreatment is required for IHC staining.

## Immunogen

The T4-positive CTL clone HG-38 was used as the immunogen for the HLA-DQ antibody.

## Storage

Store the HLA-DQ antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).