
HLA-DR Antibody Cocktail (MHC II) [clone LN3 + HLA-DRB/1067] (V2592)


Catalog No.	Formulation	Size
V2592-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2592-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2592SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2592IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	LN3 + HLA-DRB/1067
Purity	Protein G affinity chromatography
UniProt	P01911
Localization	Cell surface
Applications	Flow cytometry: 1-2ug/million cells Immunofluorescence: 1-3ug/ml Western blot: 2-4ug/ml Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This HLA-DR antibody cocktail is available for research use only.




Immunofluorescent staining of human Raji cells with HLA-DR antibody (clones LN3 + HLA-DRB/1067, cyan) and Reddot nuclear stain (red).




Western blot testing of human spleen tissue lysate with HLA-DR antibody cocktail (clones LN3 + HLA-DRB/1067). Expected molecular weight ~30 kDa.




Western blot testing of human Ramos cell lysate with HLA-DR antibody cocktail (clones LN3 + HLA-DRB/1067). Expected molecular weight ~30kDa.



IHC: Formalin-fixed, paraffin-embedded human tonsil stained with HLA-DR antibody (LN3 + HLA-DRB/1067).



IHC: Formalin-fixed, paraffin-embedded human histiocytoma stained with HLA-DR antibody (LN3 + HLA-DRB/1067).



Flow cytometry staining of human Raji cells with HLA-DR antibody; Red=isotype control, Blue= HLA-DR antibody.

Description

This mAb reacts with the HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in paraffin sections producing intense cytoplasmic staining. True histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts. Loss of HLA-DR expression is related to tumor microenvironment and predicts adverse outcome in diffuse large B-cell lymphoma.

For additional information on HLA-DR biology and complementary antibody validation, visit our HLA-DR Antibody LN3 / MHC Class II Antibody page.

Application Notes

Optimal dilution of the HLA-DR antibody cocktail should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Activated human peripheral blood mononuclear cells (LN-3 & HLA-DRB/1067) were used as the immunogen for the HLA-DR antibody cocktail.

Storage

Store the HLA-DR antibody cocktail at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.