


# HMW Cytokeratin Antibody / Basic / Type II [clone 34BE12] (V2327)


Catalog No.	Formulation	Size
V2327-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2327-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20
V2327SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100
V2327IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7

# Citations (11)


## **Bulk quote request**

Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	34BE12
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
Gene ID	3848
Localization	Cytoplasmic
Applications	Flow Cytometry: 1-2ug/10^6 cells Immunofluorescence: 1-2ug/ml Western Blot: 1-2ug/ml Immunohistochemistry (FFPE): 0.25-0.5ug/ml for 30 min at RT
Limitations	This HMW Cytokeratin antibody is available for research use only.






Immunofluorescent staining of MeOH-fixed human HeLa cells with HMW Cytokeratin antibody (clone 34BE12, green) and Reddot nuclear stain (red).



IHC staining of FFPE human prostate tissue with HMW Cytokeratin antibody (clone 34BE12). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate tissue with HMW Cytokeratin antibody (clone 34BE12). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free HMW Cytokeratin antibody (clone 34BE12) as confirmation of integrity and purity.

### **Description**

HMW Cytokeratin antibody clone 34BE12 is a monoclonal antibody specific for high molecular weight cytokeratins, a group of keratin proteins expressed in stratified epithelia. These include cytokeratins such as CK1, CK5, CK10, and CK14, which form the structural framework of squamous epithelial cells. High molecular weight cytokeratins provide resilience against mechanical stress and are reliable markers of squamous cell differentiation. NSJ Bioreagents supplies HMW Cytokeratin antibody clone 34BE12 for applications in epithelial biology, developmental studies, and cancer pathology.

HMW Cytokeratin antibody clone 34BE12 produces strong cytoplasmic staining in stratified squamous epithelia, including epidermis, esophagus, and cervix. In pathology, clone 34BE12 is frequently used to identify squamous cell carcinomas, distinguishing them from adenocarcinomas and other tumor types. Its high specificity for squamous epithelia makes it indispensable in diagnostic immunohistochemistry.

In cancer biology, HMW Cytokeratin antibody clone 34BE12 is applied to classify tumors by lineage and differentiation status. It has particular value in detecting basal and squamous differentiation in prostate and lung cancers, where it helps pathologists confirm tumor subtype and origin. Detection of high molecular weight cytokeratins also assists in evaluating

tumor grade and aggressiveness.

In cell biology and developmental research, HMW Cytokeratin antibody clone 34BE12 supports studies of epithelial maturation and keratin network assembly. High molecular weight cytokeratins are critical for maintaining epithelial integrity, and their expression patterns shift during tissue development and disease. Clone 34BE12 has been used to track these changes, helping to clarify how epithelial differentiation is regulated.

HMW Cytokeratin antibody clone 34BE12 has also been used in regenerative medicine to identify progenitor and basal cell populations. Because basal cells express high molecular weight keratins, detection with this antibody enables researchers to assess tissue renewal and repair processes.

Validated for tissue and cell-based studies, HMW Cytokeratin antibody clone 34BE12 consistently delivers robust cytoplasmic staining. It has a strong record of citation in epithelial pathology and cancer research. Alternate names include high molecular weight keratin antibody, basal cell keratin antibody, squamous marker antibody, and CK34BE12 antibody.

#### **Application Notes**

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the HMW Cytokeratin antibody to be titered up or down for optimal performance.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

### **Immunogen**

Solubilized keratin extract from human stratum corneum was used as the immunogen for this HMW Cytokeratin antibody.

#### **Storage**

Store the HMW Cytokeratin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

References (2)