


# **HOMER3 Antibody (R32981)**


Catalog No.	Formulation	Size
R32981	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide
UniProt	Q9NSC5
Localization	Cytoplasmic, cell junction
Applications	Western Blot : 0.5-1ug/ml IHC (FFPE) : 1-2ug/ml
Limitations	This HOMER3 antibody is available for research use only.




Western blot testing of 1) rat brain, 2) mouse brain, 3) human U-87MG (glioma), 4) human Jurkat and 5) human HeLa cell lysate with HOMER3 antibody at 0.5ug/ml. Predicted molecular weight ~40 kDa, can be observed at 45-47 kDa.



IHC testing of FFPE human breast cancer tissue with HOMER3 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



IHC testing of FFPE human intestinal cancer tissue with HOMER3 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



IHC testing of FFPE human lung cancer tissue with HOMER3 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.

### **Description**

Homer protein homolog 3 is a protein that in humans is encoded by the HOMER3 gene. This gene encodes a member of the HOMER family of postsynaptic density scaffolding proteins that share a similar domain structure consisting of an N-terminal Enabled/vasodilator-stimulated phosphoprotein homology 1 domain which mediates protein-protein interactions, and a carboxy-terminal coiled-coil domain and two leucine zipper motifs that are involved in self-oligomerization. The encoded protein binds numerous other proteins including group I metabotropic glutamate receptors, inositol 1,4,5-trisphosphate receptors and amyloid precursor proteins and has been implicated in diverse biological functions such as neuronal signaling, T-cell activation and trafficking of amyloid beta peptides.

#### **Application Notes**

Optimal dilution of the HOMER3 antibody should be determined by the researcher.

#### **Immunogen**

A recombinant human protein corresponding to amino acids R282-A360 was used as the immunogen for the HOMER3 antibody.

#### **Storage**

After reconstitution, the HOMER3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.