


# HSD17B1 Antibody (R30694)


Catalog No.	Formulation	Size
R30694	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
UniProt	P14061
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml Immunocytochemistry : 0.5-1ug/ml
Limitations	This HSD17B1 antibody is available for research use only.




Western blot testing of 1) human T-47D, 2) human 293T and 3) rat C6 cell lysate with HSD17B1 antibody. Predicted molecular weight  $\sim$ 35 kDa.




IHC-P: HSD17B1 antibody testing of human placenta tissue with HSD17B1 antibody. Required HIER: steam section in pH6 citrate buffer for 20 min.



ICC testing of FFPE human HeLa cells with HSD17B1 antibody. Required HIER: steam section in pH6 citrate buffer for 20 min.



IHC-P: HSD17B1 antibody testing of rat ovary tissue with HSD17B1 antibody. Required HIER: steam section in pH6 citrate buffer for 20 min.



IHC-P testing of rat liver tissue with HSD17B1 antibody. Required HIER: steam section in pH6 citrate buffer for 20 min.

### **Description**

17-beta-Hydroxysteroid dehydrogenas I, also known as 17-beta-hydroxysteroid dehydrogenase or 17-ketosteroid reductase, is an enzyme that is responsible for the interconversion of estrone(E1) and estradiol(E2) as well as the interconversion of androstenedione and testosterone. HSD17B1 gene expression is detected in 34 of 42 (81%) adenomas in all tumor subtypes; HSD17B2 mRNA is detected in 18 of 42(43%) adenomas but not in prolactinomas. Human estrogenic HSD17B1 is an NADP(H)-preferring enzyme. It possesses 11- and 4-fold higher specificity toward NADP(H) over NAD(H) for oxidation and reduction, respectively, as demonstrated by kinetic studies. All four isoforms are variably expressed in human anterior pituitary adenomas, which also show HSD17B enzyme activity, suggesting that it may play an important role in regulating the local cellular levels of estradiol.

## **Application Notes**

The stated application concentrations are suggested starting amounts. Titration of the HSD17B1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

Amino acids 29-43 (QSFKVYATLRDLKTQ-human) were used as the immunogen for this HSD17B1 antibody.

#### **Storage**

After reconstitution, the HSD17B1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.