


## HSD17B1 Antibody (RQ4894)

Catalog No.	Formulation	Size
RQ4894	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
<b>UniProt</b>	P14061
<b>Applications</b>	Western Blot : 0.5-1ug/ml
<b>Limitations</b>	This HSD17B1 antibody is available for research use only.



Western blot testing of human placenta lysate with HSD17B1 antibody at 0.5ug/ml.  
Predicted molecular weight ~35 kDa.

## Description

Estradiol 17-beta-dehydrogenase 1 is an enzyme that in humans is encoded by the HSD17B1 gene. This gene encodes a member of the 17beta-hydroxysteroid dehydrogenase family of short-chain dehydrogenases/reductases. It has a dual function in estrogen activation and androgen inactivation and plays a major role in establishing the estrogen E2 concentration gradient between serum and peripheral tissues. The encoded protein catalyzes the last step in estrogen activation, using NADPH to convert estrogens E1 and E2 and androgens like 4-androstenedione, to testosterone. It has an N-terminal short-chain dehydrogenase domain with a cofactor binding site, and a narrow, hydrophobic C-terminal domain with a steroid substrate binding site. This gene is expressed primarily in the placenta and ovarian granulosa cells, and to a lesser extent, in the endometrium, adipose tissue, and prostate. Polymorphisms in this gene have been linked to

breast and prostate cancer. A pseudogene of this gene has been identified. Alternative splicing results in multiple transcript variants.

## **Application Notes**

Optimal dilution of the HSD17B1 antibody should be determined by the researcher.

## **Immunogen**

Amino acids A20-K249 from the human protein were used as the immunogen for the HSD17B1 antibody.

## **Storage**

After reconstitution, the HSD17B1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.