


HSD17B13 Antibody / 17-beta-HSD 13 [clone HSD17B13/13106] (V5742)


Catalog No.	Formulation	Size
V5742-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5742-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5742SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	HSD17B13/13106
Purity	Protein G affinity
UniProt	Q7Z5P4
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This HSD17B13 antibody is available for research use only.



IHC staining of FFPE human hepatocellular carcinoma tissue with HSD17B13 antibody (clone HSD17B13/13106). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Description

Hydroxysteroid 17-beta dehydrogenase 13 (HSD17B13) is an enzyme in the liver that is associated with lipid droplets. It is encoded by the HSD17B13 gene in humans. HSD17B13 levels increase in patients with non-alcoholic fatty liver disease (NAFLD) and can enhance lipogenesis. However, some studies have shown that loss-of-function variants in HSD17B13 may protect against the progression of NAFLD to non-alcoholic steatohepatitis, fibrosis, and hepatocellular carcinoma.

Application Notes

Optimal dilution of the HSD17B13 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 of the human protein was used as the immunogen for the HSD17B13 antibody.

Storage

Aliquot the HSD17B13 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.