


ICAM-1 Antibody / CD54 [clone ICAM1/6606] (V4616)


Catalog No.	Formulation	Size
V4616-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4616-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4616SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	ICAM1/6606
Purity	Protein A/G affinity
UniProt	P05362
Localization	Cell Surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This ICAM-1 antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with ICAM-1 antibody (clone ICAM1/6606). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD54 / ICAM-1 antibody (clone ICAM1/6606) as confirmation of integrity and purity.

Description

Recognizes an 85-115kDa protein (variation with cell type), identified as intercellular adhesion molecule (ICAM-1). It has 7 potential N-linked glycosylation sites. ICAM-1 is a single chain glycoprotein of Ig supergene family, present on unstimulated endothelial cells (EC) and on a variety of other cell types including activated fibroblasts, EC, macrophages, and lymphocytes. ICAM-1 mediates cell adhesion by binding to integrins CD11a/CD18 (leukocyte adhesion molecule, LFA-1) and to CD11b/CD18 (Mac-1). This interaction enhances antigen-specific T-cell activation. ICAM-1 also binds to CD43 and to Plasmodium falciparum infected RBCs. ICAM-1 may also be related to progression and metastasis of tumors.

Application Notes

Optimal dilution of the ICAM-1 antibody should be determined by the researcher.

Immunogen

A recombinant fragment of the human protein was used as the immunogen for the ICAM-1 antibody.

Storage

Aliquot the ICAM-1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.