


# ICAM1 Antibody (RQ7001)


Catalog No.	Formulation	Size
RQ7001	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P05362
Applications	Western Blot : 0.5-1 ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This ICAM1 antibody is available for research use only.



Flow cytometry testing of human HepG2 cells with ICAM1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ICAM1 antibody.



Western blot testing of human 1) Ramos and 2) Raji cell lysate with ICAM1 antibody. Predicted molecular weight: ~58 kDa (unmodified), 75-115 kDa (glycosylated).

### **Description**

CD54, also known as ICAM-1 and Intercellular adhesion molecule-1 (ICAM1) is a ligand for lymphocyte function-associated (LFA) antigens. ICAM-1 is an integral membrane protein, a member of the immunoglobulin superfamily, and a ligand for LFA-1, a beta 2 leukocyte integrin. This protein is the major human rhinovirus receptor. The ICAM1 gene is mapped to human chromosome 19. In humans, lymphocyte adhesion to cells is mediated by the protein heterodimer CD11a/CD18 (Leu-CAMa, LFA-1) and its ligand CD54 (ICAM-1).

### **Application Notes**

Optimal dilution of the ICAM1 antibody should be determined by the researcher.

#### **Immunogen**

Recombinant human protein (amino acids D53-K519) was used as the immunogen for the ICAM1 antibody.

#### **Storage**

After reconstitution, the ICAM1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.