


ICAM3 Antibody [clone ICAM3/1019] (V2608)


Catalog No.	Formulation	Size
V2608-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2608-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2608SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2608IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	ICAM3/1019
Purity	Protein G affinity chromatography
UniProt	P32942
Localization	Membrane, cytoplasm
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This ICAM3 antibody is available for research use only.



Western blot analysis of HL-60 cell lysate using ICAM3 antibody (ICAM3/1019).
Observed molecular weight: 60~120 kDa depending on glycosylation level.



IHC: Formalin-fixed, paraffin-embedded human tonsil stained with ICAM3 antibody (ICAM3/1019)

Description

Recognizes an N-glycosylated glycoprotein of 120kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3. CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhesion. It is expressed on thymocytes and T lymphocytes and is resistant to treatment with phosphatidylinositol (PI) phospholipase C. This mAb is excellent for staining of formalin/paraffin tissues.

Application Notes

Optimal dilution of the ICAM3 antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the ICAM3 antibody.

Storage

Store the ICAM3 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Alternate Names

CD50