


ICOS Antibody / Inducible T-cell costimulator / CD278 [clone RM417] (R20433)

| Catalog No. | Formulation | Size |
|--------------|------------------------------------------------------------------|--------|
| R20433-0.1ML | Antibody in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide | 100 ul |


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

| | |
|---------------------------|--------------------------------------------------------------------|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Rabbit |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG |
| Clone Name | RM417 |
| Purity | Protein A purified from animal origin-free supernatant |
| UniProt | Q9Y6W8 |
| Localization | Cytoplasmic, cell membrane |
| Applications | Immunohistochemistry (FFPE) : 1:50 -1:100 |
| Limitations | This recombinant ICOS antibody is available for research use only. |



IHC staining of FFPE human appendix tissue with recombinant ICOS antibody at 1:100.



IHC staining of FFPE human tonsil tissue with recombinant ICOS antibody at 1:100.

Description

This antibody reacts to human ICOS/CD278.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant ICOS antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A peptide corresponding to the C-terminus of human Inducible T-cell costimulator protein was used as the immunogen for the recombinant ICOS antibody.

Storage

Store the recombinant ICOS antibody at -20oC.