


IFT88 Antibody (R34499)


Catalog No.	Formulation	Size
R34499-100UG	0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Antigen affinity purified
Host	Goat
Clonality	Polyclonal (goat origin)
Isotype	Goat Ig
Purity	Antigen affinity
Gene ID	8100
Localization	Cytoplasmic, cytoskeleton
Applications	Western Blot : 2-3ug/ml Immunofluorescence : 10ug/ml Flow Cytometry : 1ug/million cells in 0.1ml ELISA (peptide) LOD : 1:8000
Limitations	This IFT88 antibody is available for research use only.




Immunofluorescent staining of fixed and permeabilized human HepG2 cells with IFT88 antibody (green) at 10ug/ml and DAPI nuclear stain (blue).



Immunofluorescent staining of fixed and permeabilized human A549 cells with IFT88 antibody (green) at 10ug/ml and DAPI nuclear stain (blue).



Western blot testing of A) human K562 cell lysate without and with blocking peptide, B) human ThP-1 cell lysate without and with blocking peptide, C) human U-251 cell lysate without and with blocking peptide and D) rat KNRK cell lysate without and with blocking peptide, with IFT88 antibody.



FACS testing of fixed and permeabilized human ThP1 cells with IFT88 antibody (blue) at 1ug/million cells/0.1ml and naive goat Ig (black).

Description

Additional name(s) for this target protein: Intraflagellar transport 88, TTC10 antibody, tetratricopeptide repeat domain 10, Polaris

Application Notes

Optimal dilution of the IFT88 antibody should be determined by the researcher.

Immunogen

Amino acids KKRIDEDDFAEDEE were used as the immunogen for this IFT88 antibody.

Storage

Aliquot and store the IFT88 antibody at -20oC.