


# IGF1 Receptor Antibody / IGF1R (F54940)

Catalog No.	Formulation	Size
F54940-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54940-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	P08069
Applications	Immunohistochemistry (FFPE): 1:50-1:100 Western Blot: 1:500-1:1000
Limitations	This IGF1 Receptor antibody is available for research use only.



Western blot testing of 1) human 293T, 2) human A431, 3) mouse C2C12, 4) rat C6, 5) human HeLa and 6) mouse NIH 3T3 cell lysate with IGF1 Receptor antibody. Expected molecular weight: ~69 kDa (unglycosylated beta chain) up to ~200 kDa (glycosylated proform).

kDa
250
150 <b>=</b> 100 75
50
37 25 20

Western blot testing of human SK-BR-3 cell lysate with IGF1 Receptor antibody. Expected molecular weight: ~69 kDa (unglycosylated beta chain) up to ~200 kDa (glycosylated pro-form).



IHC testing of FFPE human cancer tissue with IGF1 Receptor antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

## **Description**

The IGF1 receptor binds insulin-like growth factor with a high affinity and plays a critical role in transformation events. Cleavage of the precursor generates alpha and beta subunits. It is highly overexpressed in most malignant tissues where it functions as an anti-apoptotic agent by enhancing cell survival. The protein possess tyrosine kinase activity.

# **Application Notes**

The stated application concentrations are suggested starting points. Titration of the IGF1 Receptor antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 1335-1366 from the human protein was used as the immunogen for the IGF1 Receptor antibody.

## **Storage**

Aliquot the IGF1 Receptor antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.