


IgM Antibody [clone IGHM/1623] (V5399)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V5399-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V5399-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V5399SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

[Bulk quote request](#)

| | |
|--------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Mouse |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG1, kappa |
| Clone Name | IGHM/1623 |
| Purity | Protein A/G affinity |
| UniProt | P01871 |
| Localization | Cell membrane, Secreted |
| Applications | ELISA : |
| Limitations | This IgM antibody is available for research use only. |



SDS-PAGE analysis of purified, BSA-free IgM antibody (clone IGHM/1623) as confirmation of integrity and purity.

Description

Recognizes a protein of 75kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes.

Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. IgM antibody is prominent in early immune responses to most antigens. Aberrant levels are associated with immune deficiency states, hereditary deficiencies, myeloma, Waldenstrom's macroglobulinemia, chronic infection and hepatocellular disease. This MAb is useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin's lymphomas. The most common feature of these malignancies is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant.

Application Notes

Optimal dilution of the IgM antibody should be determined by the researcher.

Immunogen

The heavy chain of human IgM was used as the immunogen for the IgM antibody.

Storage

Aliquot the IgM antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.