


II-10 Antibody (RQ4389)

| Catalog No. | Formulation | Size |
|-------------|---|--------|
| RQ4389 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request

| Availability | 1-3 business days |
|--------------------|---|
| Species Reactivity | Mouse |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Antigen affinity purified |
| Buffer | Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide |
| UniProt | P18893 |
| Applications | Western Blot: 0.5-1ug/ml ELISA: 0.1-0.5ug/ml (Capture; recombinant mouse protein) (BSA-free format available) |
| Limitations | This II-10 antibody is available for research use only. |



Description

Inhibits the synthesis of a number of cytokines, including IFN-gamma, IL-2, IL-3, TNF and GM-CSF produced by activated macrophages and by helper T-cells. [UniProt]

Application Notes

Optimal dilution of the II-10 antibody should be determined by the researcher.

Immunogen

A mouse partial recombinant protein corresponding to amino acids S19-S178 was used as the immunogen for the II-10 antibody.

Storage

After reconstitution, the II-10 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.