


IL-13 Antibody (R32710)


Catalog No.	Formulation	Size
R32710	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA, 0.025% sodium azide
UniProt	P35225
Localization	Cytoplasmic, membranous, secreted
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This IL-13 antibody is available for research use only.



IHC testing of FFPE human tonsil tissue with IL-13 antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



Western blot testing of human 1) HepG2 and 2) K562 cell lysate with IL-13 antibody at 0.5ug/ml. Predicted molecular weight: 13-16 kDa.

Description

Interleukin 13 is a protein that in humans is encoded by the IL-13 gene. It is a kind of cytokine secreted by many cell types, but especially T helper type 2 (Th2) cells, which is an important mediator of allergic inflammation and disease. The IL-13 gene is mapped to 5q23-q31. IL-13 induces its effects through a multi-subunit receptor that includes the alpha chain of the IL-4 receptor (IL-4RA) and at least one of two known IL-13-specific binding chains. Furthermore, this gene acts more prominently as a molecular bridge linking allergic inflammatory cells to the non-immune cells in contact with them, thereby altering physiological function.

Application Notes

Optimal dilution of the IL-13 antibody should be determined by the researcher.

Immunogen

Amino acids P36-N146 from the human protein were used as the immunogen for the IL-13 antibody.

Storage

After reconstitution, the IL-13 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.