


## IL3 Antibody (R30148)

Catalog No.	Formulation	Size
R30148	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Rat
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>Gene ID</b>	24495
<b>Applications</b>	Western Blot : 0.5-1ug/ml ELISA : 0.1-0.5ug/ml (rat protein tested); request BSA free format for coating
<b>Limitations</b>	This IL3 antibody is available for research use only.



Western blot testing of IL3 antibody and Lane 1: PC-12; 2: NRK; 3: RH35; The glycoprotein is routinely visualized between 18~35 kDa.



Western blot testing of IL3 antibody and recombinant protein (0.5ng)

## Description

Interleukin 3 is a glycoprotein that in humans is encoded by the IL3 gene. It is mapped to 5q31.1. IL-3 is an interleukin, a type of biological signal (cytokine) that can improve the body's natural response to disease as part of the immune system. It acts by binding to the interleukin-3 receptor. IL-3 stimulates the differentiation of multipotent hematopoietic stem cells into myeloid progenitor cells or, with the addition of IL-7, into lymphoid progenitor cells. In addition, IL-3 stimulates proliferation of all cells in the myeloid lineage (granulocytes, monocytes, and dendritic cells), in conjunction with other cytokines, e.g., Erythropoietin (EPO), Granulocyte macrophage colony-stimulating factor (GM-CSF), and IL-6. IL-3 is secreted by basophils and activated T cells to support growth and differentiation of T cells from the bone marrow in an immune response.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the IL3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

Rat partial recombinant protein (AA 28-166) was used as the immunogen for this IL3 antibody.

## Storage

After reconstitution, the IL3 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.