


# Insulin Antibody [clone BCPH1] (V7133)

Catalog No.	Formulation	Size
V7133-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7133-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7133SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7133IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	BCPH1
Purity	Protein G affinity chromatography
UniProt	P01308
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT (1) Prediluted IHC Only Format: incubate for 30 min at RT (2)
Limitations	This Insulin antibody is available for research use only.



IHC testing of FFPE human pancreas stained with Insulin antibody (clone BCPH1).

### **Description**

After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. [RefSeq]

## **Application Notes**

## **Immunogen**

Recombinant INS protein was used as the immunogen for the Insulin antibody.

### **Storage**

Store the Insulin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).