


Insulin Antibody [clone IRDN/805] (V2636)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V2636-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V2636-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V2636SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |
| V2636IHC-7ML | Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only* | 7 ml |

Bulk quote request

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human, Mouse |
| Format | Purified |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG1, kappa |
| Clone Name | IRDN/805 |
| Purity | Protein G affinity chromatography |
| UniProt | P01308 |
| Localization | Cytoplasmic |
| Applications | Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT |
| Limitations | This Insulin antibody is available for research use only. |



IHC: Formalin-fixed, paraffin-embedded human pancreas stained with Insulin antibody (IRDN/805).

Description

Recognizes a polypeptide which is identified as insulin, a 51-amino acid polypeptide composed of A and B chains connected through the C-peptide. Proinsulin, which has very little biological activity, is cleaved by proteases within its cell of origin into the insulin molecule and the C-terminal basic residue. Insulin enhances membrane transport of glucose, amino acids, and certain ions. It also promotes glycogen storage, formation of triglycerides, and synthesis of proteins and nucleic acids. Deficiency of insulin results in diabetes mellitus. The main storage site for insulin is the pancreatic islets. Antibodies to insulin are important as beta-cell and insulinoma marker.

Application Notes

Optimal dilution of the Insulin antibody should be determined by the researcher.

1. No special pretreatment is required for staining of formalin/paraffin tissues.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant INS protein was used as the immunogen for the Insulin antibody.

Storage

Store the Insulin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).