
Insulin Antibody IRDN/805 / Mouse Monoclonal [clone IRDN/805] (V2636)

Catalog No.	Formulation	Size
V2636-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2636-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2636SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2636IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	IRDN/805
Purity	Protein G affinity chromatography
UniProt	P01308
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Limitations	This Insulin Antibody IRDN/805 is available for research use only.



Insulin Antibody IRDN/805 IHC: Formalin-fixed, paraffin-embedded human pancreas stained with Insulin antibody (clone IRDN/805).

Description

Insulin Antibody IRDN/805 recognizes a polypeptide which is identified as insulin, a 51-amino acid polypeptide composed of A and B chains connected through the C-peptide. Proinsulin, which has very little biological activity, is cleaved by proteases within its cell of origin into the insulin molecule and the C-terminal basic residue. Insulin enhances membrane transport of glucose, amino acids, and certain ions. It also promotes glycogen storage, formation of triglycerides, and synthesis of proteins and nucleic acids. Deficiency of insulin results in diabetes mellitus. The main storage site for insulin is the pancreatic islets. Antibodies to insulin are important as beta-cell and insulinoma marker.

For a highly published insulin monoclonal with an alternate INS05 clone designation, see our Insulin Antibody 2D11-H5 / INS05 Antibody page.

Application Notes

Optimal dilution of the Insulin Antibody IRDN/805 should be determined by the researcher.

1. No special pretreatment is required for staining of formalin/paraffin tissues.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant INS protein was used as the immunogen for the Insulin Antibody IRDN/805.

Storage

Store the Insulin Antibody IRDN/805 at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.