


Interferon regulatory factor 3 Antibody / IRF3 (RQ6384)


Catalog No.	Formulation	Size
RQ6384	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q14653
Localization	Cytoplasmic, nuclear
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence (FFPE) : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This Interferon regulatory factor 3 antibody is available for research use only.




Interferon regulatory factor 3 Antibody A431 IF. Immunofluorescent staining of FFPE human A431 cells with Interferon regulatory factor 3 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human 1) HeLa, 2) Raji, 3) HepG2 and 4) mouse lung lysate with Interferon regulatory factor 3 antibody. Predicted molecular weight ~47 kDa.



Flow cytometry testing of human K562 cells with Interferon regulatory factor 3 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Interferon regulatory factor 3 antibody.



IHC staining of FFPE human rectal cancer tissue with Interferon regulatory factor 3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

IRF3 (interferon regulatory factor 3) is a member of the interferon regulatory transcription factor (IRF) family. The IRF3 gene is mapped on 19q13.33. IRF3 is found in an inactive cytoplasmic form that upon serine/threonine phosphorylation forms a complex with CREBBP. IRF3 plays an important role in the innate immune system's response to viral infection. Aggregated MAVS have been found to activate IRF3 dimerization. Although IRF3 increased transcriptional activity from an ISRE-containing promoter, expression of IRF3 as a Gal4 fusion protein did not activate expression of a chloramphenicol acetyltransferase(CAT) reporter gene containing repeats of the Gal4-binding sites. Translocation of IRF3 was accompanied by an increase in serine and threonine phosphorylation. The transcriptional activators CREBBP and EP300 coimmunoprecipitated with IRF3 only subsequent to viral infection, and the authors stated that these are also subunits of DRAF1.

For broader characterization of IRF3-associated innate immune signaling and interferon regulatory pathways, see our IRF3 Antibody / Innate Immune Signaling Antibody page featuring clone PCR-IRF3-6C8 with WB, IF, FACS, and HuProt(TM) microarray specificity validation data.

Application Notes

Optimal dilution of the Interferon regulatory factor 3 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids M1-S427) was used as the immunogen for the Interferon regulatory factor 3 antibody.

Storage

After reconstitution, the Interferon regulatory factor 3 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

