


Interleukin 17A Antibody (RQ7279)

Catalog No.	Formulation	Size
RQ7279	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q16552
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml
Limitations	This Interleukin 17A antibody is available for research use only.



IHC staining of FFPE human spleen tissue with Interleukin 17A antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

Interleukin-17A is a protein that in humans is encoded by the IL17A gene. This gene is a member of the IL-17 receptor family which includes five members (IL-17RA-E) and the encoded protein is a proinflammatory cytokine produced by activated T cells. IL-17A-mediated downstream pathways induce the production of inflammatory molecules, chemokines, antimicrobial peptides, and remodeling proteins. The encoded protein elicits crucial impacts on host defense, cell trafficking, immune modulation, and tissue repair, with a key role in the induction of innate immune defenses. This cytokine stimulates non-hematopoietic cells and promotes chemokine production thereby attracting myeloid cells to inflammatory sites. This cytokine also regulates the activities of NF-kappaB and mitogen-activated protein kinases and can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric

oxide (NO). IL-17A plays a pivotal role in various infectious diseases, inflammatory and autoimmune disorders, and cancer. High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The lung damage induced by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL17A

Application Notes

Optimal dilution of the Interleukin 17A antibody should be determined by the researcher.

Immunogen

Amino acids EDPERYPSVIWEAKCRH were used as the immunogen for the Interleukin 17A antibody.

Storage

After reconstitution, the Interleukin 17A antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.