
ISG15 Rabbit Polyclonal Antibody (F41500)

Catalog No.	Formulation	Size
F41500-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F41500-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	P05161
Localization	Cytoplasmic
Applications	Western Blot : 1:1000
Limitations	This ISG15 Rabbit Polyclonal Antibody is available for research use only.



Western blot analysis of ISG15 Rabbit Polyclonal Antibody and 293 cell lysate (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with ISG15 gene (2). Expected molecular weight: 15-17 kDa.

Description

ISG15 is secreted from monocytes in response to type I interferons and causes natural killer (NK)-cell proliferation and an augmentation of non-MCH (major histocompatibility complex)-restricted cytotoxicity. Synthesis is stimulated by IFN-alpha or IFN-beta or IFN-omega , but not IFN-gamma . ISG15 expression is also induced by overexpression of interferon regulatory factors that participate in transcriptional regulation of IFN genes, and by influenza B virus. ISG15 is secreted also by cell lines of monocyte, T-lymphocyte, B-lymphocyte, human fibroblasts, and epithelial origins. The induction of

terminal differentiation in human melanoma cells is associated with alterations in ISG15 expression. Enhancement of NK cell proliferation, augmentation of non-major histocompatibility complex-restricted cytotoxicity, and induction of IFN-gamma from T cells identify ISG15 as a member of the cytokine cascade and suggest that it may be responsible for amplifying and directing some of the immunomodulatory effects of IFN-alpha or IFN-beta. ISG15 has has also been shown to function intracellularly as a ubiquitin homolog.

For additional research into interferon-stimulated gene 15, explore our ISG15 Antibody, which provides complementary validation data for investigating ISG15 expression and interferon-responsive signaling.

Application Notes

Titration of the ISG15 Rabbit Polyclonal Antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 72-99 from the human protein was used as the immunogen for this ISG15 antibody.

Storage

Aliquot the ISG15 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Copyright © NSJ Bioreagents. All rights reserved.