
KCNA1 Antibody / Kv1.1 potassium channel (RQ6138)


Catalog No.	Formulation	Size
RQ6138	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q09470
Applications	Western Blot : 1-2ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This KCNA1 antibody is available for research use only.



Flow cytometry testing of human U-87 MG cells with KCNA1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= KCNA1 antibody.



Flow cytometry testing of human U-2 OS cells with KCNA1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= KCNA1 antibody.



Western blot testing of 1) rat brain and 2) mouse brain lysate with KCNA1 antibody. Expected molecular weight: 56-85 kDa depending on glycosylation level.

Description

Potassium voltage-gated channel subfamily A member 1, also known as Kv1.1, is a shaker related voltage-gated potassium channel that in humans is encoded by the KCNA1 gene. It is mapped to 12p13.32. The protein functions as a potassium selective channel through which the potassium ion may pass through in consensus with the electrochemical gradient. The N-terminus of the channel is associated with beta subunits that can modify the inactivation properties of the channel as well as affect expression levels. The C-terminus of the channel is complexed to a PDZ domain protein that is responsible for channel targeting.

Application Notes

Optimal dilution of the KCNA1 antibody should be determined by the researcher.

Immunogen

A human recombinant partial protein (amino acids M1-V495) was used as the immunogen for the KCNA1 antibody.

Storage

After reconstitution, the KCNA1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.