


KRT14 Antibody / CK14 / Cytokeratin 14 [clone KRT14/4125] (V9407)

Catalog No.	Formulation	Size
V9407-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9407-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9407SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2c, kappa
Clone Name	KRT14/4125
Purity	Protein A/G affinity
UniProt	P02533
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This KRT14 antibody is available for research use only.


Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using KRT14 antibody (clone KRT14/4125). These results demonstrate the foremost specificity of the KRT14/4125 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



IHC staining of FFPE human skin with KRT14 antibody (clone KRT14/4125) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free KRT14 antibody (clone KRT14/4125) as confirmation of integrity and purity.

Description

Cytokeratin 14 (CK14, KRT14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B or basic). KRT14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells.

Application Notes

Optimal dilution of the KRT14 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 351-472 was used as the immunogen for the KRT14 antibody.

Storage

Aliquot the KRT14 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.