


## LAMP-3 Antibody / CD63 [clone LAMP3/529] (V3028)


Catalog No.	Formulation	Size
V3028-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3028-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3028SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a, kappa
<b>Clone Name</b>	LAMP3/529
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P08962
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	ELISA (order BSA/sodium Azide-free Format For Coating) : Flow Cytometry : 1-2ug/million cells Immunofluorescence : 0.5-1ug/ml
<b>Limitations</b>	This LAMP-3 antibody is available for research use only.



Flow cytometry testing of PFA-fixed human U-87 MG cells with LAMP-3 antibody (clone LAMP3/529); Red=isotype control, Blue= LAMP-3 antibody.



SDS-PAGE analysis of purified, BSA-free LAMP-3 antibody as confirmation of integrity and purity.

## Description

This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63/LAMP-3. Its epitope is different from that of mAb LAMP3/803 or LAMP3/968 or NK1/C3 or MX-49.129.5. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63/LAMP-3 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. It is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63/LAMP-3 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. Deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

## Application Notes

Optimal dilution of the LAMP-3 antibody should be determined by the researcher.

## Immunogen

Recombinant human protein was used as the immunogen for the LAMP-3 antibody.

## Storage

Store the LAMP-3 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).