


## LAMP-3 Antibody / CD63 [clone LAMP3/968] (V3029)


Catalog No.	Formulation	Size
V3029-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3029-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3029SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3029IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a, kappa
<b>Clone Name</b>	LAMP3/968
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P08962
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Flow Cytometry : 1-2ug/10 <sup>6</sup> cells Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This LAMP-3 antibody is available for research use only.




Flow cytometry testing of human MCF7 cells with LAMP-3 antibody (clone LAMP3/968); Gray=unstained, Orange= CF555-labeled LAMP-3 antibody.




FACS staining of bead-bound exosomes derived from human MCF7 cells: Gray = unstained, Orange = CF555-labeled LAMP-3 antibody.




Flow cytometry testing of PFA-fixed human U-87 MG cells with LAMP-3 antibody (clone LAMP3/968); Red=isotype control, Blue= LAMP-3 antibody.



IHC: Formalin-fixed, paraffin-embedded human prostate carcinoma stained with CD63 / LAMP-3 antibody (LAMP3/968).



IHC: Formalin-fixed, paraffin-embedded human melanoma stained with CD63 / LAMP-3 antibody (LAMP3/968).



SDS-PAGE Analysis of Purified, BSA-Free LAMP-3 Antibody (clone LAMP3/968). Confirmation of Integrity and Purity of the Antibody.

## Description

This mAb recognizes protein of 26kDa-60kDa, which is identified as CD63/LAMP-3. Its epitope is different from that of mAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63/LAMP-3 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. It is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53,

which contain four transmembrane regions. CD63/LAMP-3 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. Deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

## Application Notes

Optimal dilution of the LAMP-3 antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

Recombinant human full-length protein was used as the immunogen for the CD63 / LAMP-3 antibody.

## Storage

Store the LAMP-3 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).