


## LCAT Antibody (R31981)

Catalog No.	Formulation	Size
R31981	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>UniProt</b>	P16301
<b>Applications</b>	Western Blot : 0.1-0.5ug/ml
<b>Limitations</b>	This LCAT antibody is available for research use only.



Western blot testing of 1) rat brain, 2) mouse testis, 3) human HepG2 and 4) human HeLa lysate with LCAT antibody. Observed molecular weight: 50-75 kDa depending on glycosylation level.

## Description

LCAT (Lecithin: Cholesterol Acyltransferase), is an enzyme that converts free cholesterol into cholesteryl ester. Azoulay et al. (1987) used a cDNA clone corresponding to LCAT to assign the locus to 16q22 through the analysis of DNA from somatic cell hybrids and in situ hybridization. LCAT plays an important role in lipoprotein metabolism, especially in the process termed 'reverse cholesterol transport.' The enzyme is synthesized in the liver and circulates in blood plasma as a complex with components of high density lipoprotein (HDL). Cholesterol from peripheral cells is transferred to HDL particles, esterified through the action of LCAT on HDL, and incorporated into the core of the lipoprotein. The cholesterol ester is thereby transported to the liver (Jonas, 2000).

## Application Notes

Optimal dilution of the LCAT antibody should be determined by the researcher.

## Immunogen

Amino acids QPVHLLPMNETDHLNMVFSNKTLEHINAILLGAYR of mouse LCAT were used as the immunogen for the LCAT antibody.

## Storage

After reconstitution, the LCAT antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.