


## LCK Antibody [clone 845CT3.5.2] (F53783)

Catalog No.	Formulation	Size
F53783-0.1ML	In ascites with 0.09% sodium azide	0.1 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Ascites
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG1
<b>Clone Name</b>	845CT3.5.2
<b>Purity</b>	Ascites
<b>UniProt</b>	P06239
<b>Applications</b>	Western Blot : 1:5000
<b>Limitations</b>	This LCK antibody is available for research use only.



LCK antibody western blot analysis in 1) HL-60, 2) Jurkat and 3) NCI-H292 lysate. Predicted molecular weight ~58 kDa.

### Description

Non-receptor tyrosine-protein kinase that plays an essential role in the selection and maturation of developing T-cells in the thymus and in the function of mature T-cells. Plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways. Constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors. Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, thereby recruiting the associated LCK protein to the vicinity of the TCR/CD3 complex. LCK then phosphorylates tyrosines residues within the immunoreceptor tyrosine-based activation motifs (ITAM) of the cytoplasmic tails of the TCR-gamma chains and CD3 subunits, initiating the TCR/CD3 signaling pathway. Once

stimulated, the TCR recruits the tyrosine kinase ZAP70, that becomes phosphorylated and activated by LCK. Following this, a large number of signaling molecules are recruited, ultimately leading to lymphokine production. LCK also contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, which leads to hyperphosphorylation and activation of LCK. Also plays a role in the IL2 receptor-linked signaling pathway that controls the T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Phosphorylates other substrates including RUNX3, PTK2B/PYK2, the microtubule-associated protein MAPT, RHOH or TYROBP.

Learn more about LCK function in T-cell receptor signaling, lymphocyte activation, adaptive immunity, and immune signal transduction on our [LCK Antibody](#) page.

## Application Notes

Titration of the LCK antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

Purified His-tagged protein was used to produced this monoclonal LCK antibody.

## Storage

Aliquot the LCK antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.