
LIR-1 Antibody / LILRB1 / ILT-2 (RQ6492)


Catalog No.	Formulation	Size
RQ6492	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q8NHL6
Applications	Western blot: 1-2ug/ml Immunohistochemistry (FFPE): 2-5ug/ml Flow cytometry: 1-3ug/million cells Direct ELISA: 0.1-0.5ug/ml
Limitations	This LIR-1 antibody is available for research use only.




IHC staining of FFPE human lymphoma tissue with LIR-1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver cancer tissue with LIR-1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human ThP-1 and 2) human Raji cell lysate with LIR-1 antibody. Predicted molecular weight ~71 kDa.



Flow cytometry testing of human HL60 cells with LIR-1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= LIR-1 antibody.

Description

Leukocyte immunoglobulin-like receptor subfamily B member 1 is a protein that in humans is encoded by the LILRB1 gene. This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene.

Application Notes

Optimal dilution of the LIR-1 antibody should be determined by the researcher.

Immunogen

An E. coli-derived human protein (amino acids Q347-Q442) was used as the immunogen for the LIR-1 antibody.

Storage

After reconstitution, the LIR-1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.