


## LMO2 Antibody / Rhombotin 2 (F43518)

Catalog No.	Formulation	Size
F43518-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F43518-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Predicted Reactivity</b>	Mouse
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	P25791
<b>Applications</b>	Western Blot : 1:1000 IHC (Paraffin) : 1:10-1:50
<b>Limitations</b>	This LMO2 antibody is available for research use only.



LMO2 antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue.



LMO2 antibody western blot analysis in NCI-H460 lysate.

## Description

LMO2 encodes a cysteine-rich, two LIM-domain protein that is required for yolk sac erythropoiesis. The LMO2 protein has a central and crucial role in hematopoietic development and is highly conserved. The LMO2 transcription start site is located approximately 25 kb downstream from the 11p13 T-cell translocation cluster (11p13 ttc), where a number T-cell acute lymphoblastic leukemia-specific translocations occur. Alternative splicing results in multiple transcript variants encoding different isoforms.

## Application Notes

Titration of the LMO2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 1-30 from the human protein was used as the immunogen for this LMO2 antibody.

## Storage

Aliquot the LMO2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.