


# LYN Antibody [clone 1174CT22.3.1.1] (F52444)

Catalog No.	Formulation	Size
F52444-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F52444-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, k
Clone Name	1174CT22.3.1.1
Purity	Purified
UniProt	P07948
Localization	Cytoplasmic, nuclear, cell membrane
Applications	Western Blot: 1:1000
Limitations	This LYN antibody is available for research use only.



Western blot analysis of lysate from 1) A431 and 2) Raji cell line using LYN antibody. Predicted molecular weight ~53 kDa (p53lyn), ~56 kDa (p56lyn).

## **Description**

Non-receptor tyrosine-protein kinase that transmits signals from cell surface receptors and plays an important role in the regulation of innate and adaptive immune responses, hematopoiesis, responses to growth factors and cytokines, integrin signaling, but also responses to DNA damage and genotoxic agents. Functions primarily as negative regulator, but can also function as activator, depending on the context. Required for the initiation of the B-cell response, but also for its down- regulation and termination. Plays an important role in the regulation of B-cell differentiation, proliferation, survival

and apoptosis, and is important for immune self-tolerance. Acts downstream of several immune receptors, including the Bcell receptor, CD79A, CD79B, CD5, CD19, CD22, FCER1, FCGR2, FCGR1A, TLR2 and TLR4. Plays a role in the inflammatory response to bacterial lipopolysaccharide. Mediates the responses to cytokines and growth factors in hematopoietic progenitors, platelets, erythrocytes, and in mature myeloid cells, such as dendritic cells, neutrophils and eosinophils. Acts downstream of EPOR, KIT, MPL, the chemokine receptor CXCR4, as well as the receptors for IL3, IL5 and CSF2. Plays an important role in integrin signaling. Regulates cell proliferation, survival, differentiation, migration, adhesion, degranulation, and cytokine release. Down- regulates signaling pathways by phosphorylation of immunoreceptor tyrosine-based inhibitory motifs (ITIM), that then serve as binding sites for phosphatases, such as PTPN6/SHP-1, PTPN11/SHP-2 and INPP5D/SHIP-1, that modulate signaling by dephosphorylation of kinases and their substrates. Phosphorylates LIME1 in response to CD22 activation. Phosphorylates BTK, CBL, CD5, CD19, CD72, CD79A, CD79B, CSF2RB, DOK1, HCLS1, LILRB3/PIR-B, MS4A2/FCER1B, PTK2B/PYK2, SYK and TEC. Promotes phosphorylation of SIRPA, PTPN6/SHP-1, PTPN11/SHP-2 and INPP5D/SHIP-1. Mediates phosphorylation of the BCR-ABL fusion protein. Required for rapid phosphorylation of FER in response to FCER1 activation. Mediates KIT phosphorylation. Acts as an effector of EPOR (erythropoietin receptor) in controlling KIT expression and may play a role in erythroid differentiation during the switch between proliferation and maturation. Depending on the context, activates or inhibits several signaling cascades. Regulates phosphatidylinositol 3- kinase activity and AKT1 activation. Regulates activation of the MAP kinase signaling cascade, including activation of MAP2K1/MEK1, MAPK1/ERK2, MAPK3/ERK1, MAPK8/JNK1 and MAPK9/JNK2. Mediates activation of STAT5A and/or STAT5B. Phosphorylates LPXN on 'Tyr-72'.

#### **Application Notes**

Titration of the LYN antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A recombinant protein corresponding to amino acids 1-210 from the human protein was used as the immunogen for the LYN antibody.

#### **Storage**

Aliquot the LYN antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.