


## Mannose-binding protein C Antibody / MBL2 (RQ8126)


Catalog No.	Formulation	Size
RQ8126	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	P11226
<b>Applications</b>	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This Mannose-binding protein C antibody is available for research use only.



Flow cytometry testing of fixed and permeabilized human HepG2 cells with Mannose-binding protein C antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Mannose-binding protein C antibody.



Western blot testing of human HCCT cell lysate with Mannose-binding protein C antibody. Predicted molecular weight ~26 kDa.

## Description

This gene encodes the soluble mannose-binding lectin or mannose-binding protein found in serum. The protein encoded belongs to the collectin family and is an important element in the innate immune system. The protein recognizes and binds to mannose and N-acetylglucosamine on many microorganisms, including bacteria, yeast, and viruses including influenza virus, HIV and SARS-CoV. This binding activates the classical complement pathway. Deficiencies of this gene have been associated with susceptibility to autoimmune and infectious diseases.

## Application Notes

Optimal dilution of the Mannose-binding protein C antibody should be determined by the researcher.

## Immunogen

E. coli-derived recombinant human protein (amino acids E21-I248) was used as the immunogen for the Mannose-binding protein C antibody.

## Storage

After reconstitution, the Mannose-binding protein C antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.