


## MAX Antibody / MYC associated factor X (R33757)

Catalog No.	Formulation	Size
R33757-100UG	0.5 mg/ml in 1X TBS, pH7.3, with 0.5% BSA (US sourced) and 0.02% sodium azide	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Predicted Reactivity</b>	Cow, Dog
<b>Format</b>	Antigen affinity purified
<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal (goat origin)
<b>Isotype</b>	Goat Ig
<b>Purity</b>	Antigen affinity
<b>Gene ID</b>	4149
<b>Applications</b>	Western Blot : 0.1-0.3ug/ml ELISA (peptide) LOD : 1:32000
<b>Limitations</b>	This MAX antibody is available for research use only.



Western blot testing of Jurkat lysate with MAX antibody at 0.1 ug/ml. Two forms of MAX may be observed: 16-17kDa and 21-22kDa.

### Description

Additional name(s) for this target protein: MYC associated factor X

### Application Notes

Optimal dilution of the MAX antibody should be determined by the researcher.

## Immunogen

Amino acids EEPQSRKKLRMEAS were used as the immunogen for this MAX antibody.

## Storage

Aliquot and store the MAX antibody at -20oC.

## References (1)